# ALGEBRA QUALIFYING EXAM PROBLEMS RING THEORY

Kent State University Department of Mathematical Sciences

> Compiled and Maintained by Donald L. White

Version: August 29, 2017

# CONTENTS

RING THEORY
General Ring Theory 1
Prime, Maximal, and Primary Ideals 4
Commutative Rings
Domains
Polynomial Rings
Non-commutative Rings 10
Local Rings, Localization, Rings of Fractions 11
Chains and Chain Conditions 12

# **RING THEORY**

## General Ring Theory

- 1. Give an example of each of the following.
  - (a) An irreducible polynomial of degree 3 in  $\mathbb{Z}_3[x]$ .
  - (b) A polynomial in  $\mathbb{Z}[x]$  that is not irreducible in  $\mathbb{Z}[x]$  but is irreducible in  $\mathbb{Q}[x]$ .
  - (c) A non-commutative ring of characteristic p, p a prime.
  - (d) A ring with exactly 6 invertible elements.
  - (e) An infinite non-commutative ring with only finitely many ideals.
  - (f) An infinite non-commutative ring with non-zero characteristic.
  - (g) An integral domain which is not a unique factorization domain.
  - (h) A unique factorization domain that is not a principal ideal domain.
  - (i) A principal ideal domain that is not a Euclidean domain.
  - (j) A Euclidean domain other than the ring of integers or a field.
  - (k) A finite non-commutative ring.
  - (1) A commutative ring with a sequence  $\{P_n\}_{n=1}^{\infty}$  of prime ideals such that  $P_n$  is properly contained in  $P_{n+1}$  for all n.
  - (m) A non-zero prime ideal of a commutative ring that is not a maximal ideal.
  - (n) An irreducible element of a commutative ring that is not a prime element.
  - (o) An irreducible element of an integral domain that is not a prime element.
  - (p) A commutative ring that has exactly one maximal ideal and is not a field.
  - (q) A non-commutative ring with exactly two maximal ideals.
- 2. (a) How many units does the ring  $\mathbb{Z}/60\mathbb{Z}$  have? Explain your answer.
  - (b) How many ideals does the ring  $\mathbb{Z}/60\mathbb{Z}$  have? Explain your answer.
- 3. **[NEW]** How many ideals does the ring  $\mathbb{Z}/90\mathbb{Z}$  have? Explain your answer.
- 4. Denote the set of invertible elements of the ring  $\mathbb{Z}_n$  by  $U_n$ .
  - (a) List all the elements of  $U_{18}$ .
  - (b) Is  $U_{18}$  a cyclic group under multiplication? Justify your answer.
- 5. **[NEW]** Denote the set of invertible elements of the ring  $\mathbb{Z}_n$  by  $U_n$ .
  - (a) List all the elements of  $U_{24}$ .
  - (b) Is  $U_{24}$  a cyclic group under multiplication? Justify your answer.
- 6. [NEW] Find all positive integers n having the property that the group of units of  $\mathbb{Z}/n\mathbb{Z}$  is an elementary abelian 2-group.
- 7. Let U(R) denote the group of units of a ring R. Prove that if m divides n, then the natural ring homomorphism  $\mathbb{Z}_n \to \mathbb{Z}_m$  maps  $U(\mathbb{Z}_n)$  onto  $U(\mathbb{Z}_m)$ . Give an example that shows that U(R) does not have to map onto U(S) under a surjective ring homomorphism  $R \to S$ .

- 8. If p is a prime satisfying  $p \equiv 1 \pmod{4}$ , then p is a sum of two squares.
- 9. If  $(\frac{1}{2})$  denotes the Legendre symbol, prove Euler's Criterion: if p is a prime and a is any integer relatively prime to p, then  $a^{(p-1)/2} \equiv \left(\frac{a}{p}\right) \pmod{p}$ .
- 10. Let  $R_1$  and  $R_2$  be commutative rings with identities and let  $R = R_1 \times R_2$ . Show that every ideal I of R is of the form  $I = I_1 \times I_2$  with  $I_i$  an ideal of  $R_i$  for i = 1, 2.
- 11. Show that a non-zero ring R in which  $x^2 = x$  for all  $x \in R$  is of characteristic 2 and is commutative.
- 12. Let R be a finite commutative ring with more than one element and no zero-divisors. Show that R is a field.
- 13. Determine for which integers n the ring  $\mathbb{Z}/n\mathbb{Z}$  is a direct sum of fields. Prove your answer.
- 14. Let R be a subring of a field F such that for each x in F either  $x \in R$  or  $x^{-1} \in R$ . Prove that if I and J are two ideals of R, then either  $I \subseteq J$  or  $J \subseteq I$ .
- 15. The Jacobson Radical J(R) of a ring R is defined to be the intersection of all maximal ideals of R. Let R be a commutative ring with 1 and let  $x \in R$ . Show that  $x \in J(R)$  if and only if 1 - xy

Let *R* be a commutative ring with 1 and let  $x \in R$ . Show that  $x \in J(R)$  if and only if 1 - xy is a unit for all *y* in *R*.

- 16. Let R be any ring with identity, and n any positive integer. If  $M_n(R)$  denotes the ring of  $n \times n$  matrices with entries in R, prove that  $M_n(I)$  is an ideal of  $M_n(R)$  whenever I is an ideal of R, and that every ideal of  $M_n(R)$  has this form.
- 17. Let m, n be positive integers such that m divides n. Then the natural map  $\varphi : \mathbb{Z}_n \to \mathbb{Z}_m$  given by  $a + (n) \mapsto a + (m)$  is a surjective ring homomorphism. If  $U_n, U_m$  are the units of  $\mathbb{Z}_n$  and  $\mathbb{Z}_m$ , respectively, show that  $\varphi : U_n \to U_m$  is a surjective group homomorphism.
- 18. Let R be a ring with ideals A and B. Let  $R/A \times R/B$  be the ring with coordinate-wise addition and multiplication. Show the following.
  - (a) The map  $R \to R/A \times R/B$  given by  $r \mapsto (r + A, r + B)$  is a ring homomorphism.
  - (b) The homomorphism in part (a) is surjective if and only if A + B = R.
- 19. Let m and n be relatively prime integers.
  - (a) Show that if c and d are any integers, then there is an integer x such that  $x \equiv c \pmod{m}$  and  $x \equiv d \pmod{n}$ .
  - (b) Show that  $\mathbb{Z}_{mn}$  and  $\mathbb{Z}_m \times \mathbb{Z}_n$  are isomorphic as rings.
- 20. Let R be a commutative ring with 1 and let I and J be ideals of R such that I + J = R. Show that  $R/(I \cap J) \cong R/I \oplus R/J$ .
- 21. **[NEW]** Let R be a commutative ring with identity and let  $I_1, I_2, \ldots, I_n$  be pairwise comaximal ideals of R (i.e.,  $I_i + I_j = R$  if  $i \neq j$ ). Show that  $I_i + \bigcap_{i \neq j} I_j = R$  for all i.
- 22. Let R be a commutative ring, not necessarily with identity, and assume there is some fixed positive integer n such that nr = 0 for all  $r \in R$ . Prove that R embeds in a ring S with identity so that R is an ideal of S and  $S/R \cong \mathbb{Z}/n\mathbb{Z}$ .

- 23. Let R be a ring with identity 1 and  $a, b \in R$  such that ab = 1. Denote  $X = \{x \in R \mid ax = 1\}$ . Show the following.
  - (a) If  $x \in X$ , then  $b + (1 xa) \in X$ .
  - (b) If  $\varphi: X \to X$  is the mapping given by  $\varphi(x) = b + (1 xa)$ , then  $\varphi$  is one-to-one.
  - (c) If X has more than one element, then X is an infinite set.
- 24. Let R be a commutative ring with identity and define  $U_2(R) = \left\{ \begin{bmatrix} a & b \\ 0 & c \end{bmatrix} \mid a, b, c \in R \right\}$ . Prove that every R-automorphism of  $U_2(R)$  is inner.
- 25. Let  $\mathbb{R}$  be the field of real numbers and let F be the set of all  $2 \times 2$  matrices of the form  $\begin{bmatrix} a & b \\ -3b & a \end{bmatrix}$ , where  $a, b \in \mathbb{R}$ . Show that F is a field under the usual matrix operations.

26. Let R be the ring of all  $2 \times 2$  matrices of the form  $\begin{bmatrix} a & b \\ -b & a \end{bmatrix}$  where a and b are real numbers. Prove that R is isomorphic to  $\mathbb{C}$ , the field of complex numbers.

27. Let p be a prime and let R be the ring of all  $2 \times 2$  matrices of the form  $\begin{bmatrix} a & b \\ pb & a \end{bmatrix}$ , where  $a, b \in \mathbb{Z}$ . Prove that R is isomorphic to  $\mathbb{Z}[\sqrt{p}]$ .

28. Let p be a prime and  $F_p$  the set of all  $2 \times 2$  matrices of the form  $\begin{bmatrix} a & b \\ -b & a \end{bmatrix}$ , where  $a, b \in \mathbb{Z}_p$ .

- (a) Show that  $F_p$  is a commutative ring with identity.
- (b) Show that  $F_7$  is a field.
- (c) Show that  $F_{13}$  is not a field.
- 29. Let  $I \subseteq J$  be right ideals of a ring R such that  $J/I \cong R$  as right R-modules. Prove that there exists a right ideal K such that  $I \cap K = (0)$  and I + K = J.
- 30. A ring R is called simple if  $R^2 \neq 0$  and 0 and R are its only ideals. Show that the center of a simple ring is 0 or a field.
- 31. Give an example of a field F and a one-to-one ring homomorphism  $\varphi: F \to F$  which is not onto. Verify your example.
- 32. Let D be an integral domain and let  $D[x_1, x_2, \ldots, x_n]$  be the polynomial ring over D in the n indeterminates  $x_1, x_2, \ldots, x_n$ . Let

$$V = \begin{bmatrix} x_1^{n-1} & \cdots & x_1^2 & x_1 & 1 \\ x_2^{n-1} & \cdots & x_2^2 & x_2 & 1 \\ \vdots & & \vdots & \vdots & \vdots \\ x_n^{n-1} & \cdots & x_n^2 & x_n & 1 \end{bmatrix}$$

Prove that the determinant of V is  $\prod_{1 \leq i < j \leq n} (x_i - x_j).$ 

33. Let R = C[0, 1] be the set of all continuous real-valued functions on [0, 1]. Define addition and multiplication on R as follows. For  $f, g \in R$  and  $x \in [0, 1]$ ,

$$(f+g)(x) = f(x) + g(x)$$
 and  $(fg)(x) = f(x)g(x)$ .

- (a) Show that R with these operations is a commutative ring with identity.
- (b) Find the units of R.
- (c) If  $f \in R$  and  $f^2 = f$ , then  $f = 0_R$  or  $f = 1_R$ .
- (d) If n is a positive integer and  $f \in R$  is such that  $f^n = 0_R$ , then  $f = 0_R$ .
- 34. Let S be the ring of all bounded, continuous functions  $f : \mathbb{R} \to \mathbb{R}$ , where  $\mathbb{R}$  is the set of real numbers. Let I be the set of functions f in S such that  $f(t) \to 0$  as  $|t| \to \infty$ .
  - (a) Show that I is an ideal of S.
  - (b) Suppose  $x \in S$  is such that there is an  $i \in I$  with ix = x. Show that x(t) = 0 for all sufficiently large |t|.
- 35. Let  $\mathbb{Q}$  be the field of rational numbers and  $D = \{a + b\sqrt{2} \mid a, b \in \mathbb{Q}\}.$ 
  - (a) Show that D is a subring of the field of real numbers.
  - (b) Show that D is a principal ideal domain.
  - (c) Show that  $\sqrt{3}$  is not an element of D.
- 36. Show that if p is a prime such that  $p \equiv 1 \pmod{4}$ , then  $x^2 + 1$  is not irreducible in  $\mathbb{Z}_p[x]$ .
- 37. Show that if p is a prime such that  $p \equiv 3 \pmod{4}$ , then  $x^2 + 1$  is irreducible in  $\mathbb{Z}_p[x]$ .
- 38. Show that if p is a prime such that  $p \equiv 1 \pmod{6}$ , then  $x^3 + 1$  splits in  $\mathbb{Z}_p[x]$ .

#### Prime, Maximal, and Primary Ideals

- 39. Let R be a non-zero commutative ring with 1. Show that an ideal M of R is maximal if and only if R/M is a field.
- 40. Let R be a commutative ring with 1. Show that an ideal P of R is prime if and only if R/P is an integral domain.
- 41. (a) Let R be a commutative ring with 1. Show that if M is a maximal ideal of R then M is a prime ideal of R.
  - (b) Give an example of a non-zero prime ideal in a ring R that is not a maximal ideal.
- 42. Let R be a non-zero ring with identity. Show that every proper ideal of R is contained in a maximal ideal.
- 43. **[NEW]** Let  $M_1 \neq M_2$  be two maximal ideals in the commutative ring R and let  $I = M_1 \cap M_2$ . Prove that R/I is isomorphic to the direct sum of two fields.
- 44. Let R be a non-zero commutative ring with 1. Show that if I is an ideal of R such that 1 + a is a unit in R for all  $a \in I$ , then I is contained in every maximal ideal of R.

- 45. **[NEW]** Let R be a commutative ring with identity. Suppose R contains an idempotent element a other than 0 or 1. Show that every prime ideal in R contains an idempotent element other than 0 or 1. (An element  $a \in R$  is idempotent if  $a^2 = a$ .)
- 46. Let R be a commutative ring with 1.
  - (a) Prove that (x) is a prime ideal in R[x] if and only if R is an integral domain.
  - (b) Prove that (x) is a maximal ideal in R[x] if and only if R is a field.
- 47. Find all values of a in  $\mathbb{Z}_3$  such that the quotient ring

$$\mathbb{Z}_3[x]/(x^3+x^2+ax+1)$$

is a field. Justify your answer.

48. Find all values of a in  $\mathbb{Z}_5$  such that the quotient ring

$$\mathbb{Z}_{5}[x]/(x^{3}+2x^{2}+ax+3)$$

is a field. Justify your answer.

- 49. Let R be a commutative ring with identity and let U be maximal among non-finitely generated ideals of R. Prove U is a prime ideal.
- 50. Let R be a commutative ring with identity and let U be maximal among non-principal ideals of R. Prove U is a prime ideal.
- 51. Let R be a non-zero commutative ring with 1 and S a multiplicative subset of R not containing 0. Show that if P is maximal in the set of ideals of R not intersecting S, then P is a prime ideal.
- 52. Let R be a non-zero commutative ring with 1.
  - (a) Let S be a multiplicative subset of R not containing 0 and let P be maximal in the set of ideals of R not intersecting S. Show that P is a prime ideal.
  - (b) Show that the set of nilpotent elements of R is the intersection of all prime ideals.
- 53. Let R be a commutative ring with identity and let  $x \in R$  be a non-nilpotent element. Prove that there exists a prime ideal P of R such that  $x \notin P$ .
- 54. Let R be a commutative ring with identity and let S be the set of all elements of R that are not zero-divisors. Show that there is a prime ideal P such that  $P \cap S$  is empty. (Hint: Use Zorn's Lemma.)
- 55. Let R be a commutative ring with identity and let C be a chain of prime ideals of R. Show that  $\bigcup_{P \in \mathcal{C}} P$  and  $\bigcap_{P \in \mathcal{C}} P$  are prime ideals of R.
- 56. Let R be a commutative ring and P a prime ideal of R. Show that there is a prime ideal  $P_0 \subseteq P$  that does not properly contain any prime ideal.
- 57. Let R be a commutative ring with 1 such that for every x in R there is an integer n > 1 (depending on x) such that  $x^n = x$ . Show that every prime ideal of R is maximal.
- 58. Let R be a commutative ring with 1 in which every ideal is a prime ideal. Prove that R is a field. (Hint: For  $a \neq 0$  consider the ideals (a) and ( $a^2$ ).)

- 59. Let D be a principal ideal domain. Prove that every nonzero prime ideal of D is a maximal ideal.
- 60. Show that if R is a finite commutative ring with identity then every prime ideal of R is a maximal ideal.
- 61. Let R = C[0, 1] be the ring of all continuous real-valued functions on [0, 1], with addition and multiplication defined as follows. For  $f, g \in R$  and  $x \in [0, 1]$ ,

$$(f+g)(x) = f(x) + g(x)$$
  
(fg)(x) = f(x)g(x).

Prove that if M is a maximal ideal of R, then there is a real number  $x_0 \in [0, 1]$  such that  $M = \{f \in R \mid f(x_0) = 0\}.$ 

- 62. Let R be a commutative ring with identity, and let  $P \subset Q$  be prime ideals of R. Prove that there exist prime ideals  $P^*, Q^*$  satisfying  $P \subseteq P^* \subset Q^* \subseteq Q$ , such that there are no prime ideals strictly between  $P^*$  and  $Q^*$ . HINT: Fix  $x \in Q - P$  and show that there exists a prime ideal  $P^*$  containing P, contained in Q and maximal with respect to not containing x.
- 63. Let R be a commutative ring with 1. An ideal I of R is called a *primary* ideal if  $I \neq R$  and for all  $x, y \in R$  with  $xy \in I$ , either  $x \in I$  or  $y^n \in I$  for some integer  $n \ge 1$ .
  - (a) Show that an ideal I of R is primary if and only if  $R/I \neq 0$  and every zero-divisor in R/I is nilpotent.
  - (b) Show that if I is a primary ideal of R then the radical  $\operatorname{Rad}(I)$  of I is a prime ideal. (Recall that  $\operatorname{Rad}(I) = \{x \in R \mid x^n \in I \text{ for some } n\}$ .)

# **Commutative Rings**

- 64. Let R be a commutative ring with identity. Show that R is an integral domain if and only if R is a subring of a field.
- 65. Let R be a commutative ring with identity. Show that if x and y are nilpotent elements of R then x + y is nilpotent and the set of all nilpotent elements is an ideal in R.
- 66. Let R be a commutative ring with identity. An ideal I of R is *irreducible* if it cannot be expressed as the intersection of two ideals of R neither of which is contained in the other. Show the following.
  - (a) If P is a prime ideal then P is irreducible.
  - (b) If x is a non-zero element of R, then there is an ideal  $I_x$ , maximal with respect to the property that  $x \notin I_x$ , and  $I_x$  is irreducible.
  - (c) If every irreducible ideal of R is a prime ideal, then 0 is the only nilpotent element of R.
- 67. Let R be a commutative ring with 1 and let I be an ideal of R satisfying  $I^2 = \{0\}$ . Show that if  $a + I \in R/I$  is an idempotent element of R/I, then the cos t a + I contains an idempotent element of R.
- 68. Let R be a commutative ring with identity that has exactly one prime ideal P. Prove the following.
  - (a) R/P is a field.
  - (b) R is isomorphic to  $R_P$ , the ring of quotients of R with respect to the multiplicative set  $R P = \{s \in R \mid s \notin P\}.$

- 69. Let R be a commutative ring with identity and  $\sigma: R \to R$  a ring automorphism.
  - (a) Show that  $F = \{r \in R \mid \sigma(r) = r\}$  is a subring of R and the identity of R is in F.
  - (b) Show that if  $\sigma^2$  is the identity map on R, then each element of R is the root of a monic polynomial of degree two in F[x].
- 70. Let R be a commutative ring with identity that has exactly three ideals,  $\{0\}$ , I, and R.
  - (a) Show that if  $a \notin I$ , then a is a unit of R.
  - (b) Show that if  $a, b \in I$  then ab = 0.
- 71. Let R be a commutative ring with 1. Show that if u is a unit in R and n is nilpotent, then u + n is a unit.
- 72. Let R be a commutative ring with identity. Suppose that for every  $a \in R$ , either a or 1 a is invertible. Prove that  $N = \{a \in R \mid a \text{ is not invertible}\}$  is an ideal of R.
- 73. Let R be a commutative ring with 1. Show that the sum of any two principal ideals of R is principal if and only if every finitely generated ideal of R is principal.
- 74. Let R be a commutative ring with identity such that not every ideal is a principal ideal.
  - (a) Show that there is an ideal I maximal with respect to the property that I is not a principal ideal.
  - (b) If I is the ideal of part (a), show that R/I is a principal ideal ring.
- 75. Recall that if  $R \subseteq S$  is an inclusion of commutative rings (with the same identity) then an element  $s \in S$  is *integral over* R if s satisfies some monic polynomial with coefficients in R. Prove the equivalence of the following statements.
  - (i) s is integral over R.
  - (ii) R[s] is finitely generated as an *R*-module.
  - (iii) There exists a faithful R[s] module which is finitely generated as an R-module.
- 76. Recall that if  $R \subseteq S$  is an inclusion of commutative rings (with the same identity) then S is an *integral* extension of R if every element of S satisfies some monic polynomial with coefficients in R. Prove that if  $R \subseteq S \subseteq T$  are commutative rings with the same identity, then S is integral over R and T is integral over S if and only if T is integral over R.
- 77. Let  $R \subseteq S$  be commutative domains with the same identity, and assume that S is an integral extension of R. Let I be a nonzero ideal of S. Prove that  $I \cap R$  is a nonzero ideal of R.

#### Domains

78. Suppose R is a domain and I and J are ideals of R such that IJ is principal. Show that I (and by symmetry J) is finitely generated.

[Hint: If IJ = (a), then  $a = \sum_{i=1}^{n} x_i y_i$  for some  $x_i \in I$  and  $y_i \in J$ . Show the  $x_i$  generate I.]

- 79. [NEW] Prove that if D is a Euclidean Domain, then D is a Principal Ideal Domain.
- 80. Show that if p is a prime such that there is an integer b with  $p = b^2 + 4$ , then  $\mathbb{Z}[\sqrt{p}]$  is not a unique factorization domain.

- 81. Show that if p is a prime such that  $p \equiv 1 \pmod{4}$ , then  $\mathbb{Z}[\sqrt{p}]$  is not a unique factorization domain.
- 82. Let  $D = \mathbb{Z}(\sqrt{5}) = \{m + n\sqrt{5} \mid m, n \in \mathbb{Z}\}$  a subring of the field of real numbers and necessarily an integral domain (you need not show this) and  $F = \mathbb{Q}(\sqrt{5})$  its field of fractions. Show the following:
  - (a)  $x^2 + x 1$  is irreducible in D[x] but not in F[x].
  - (b) D is not a unique factorization domain.
- 83. Let  $D = \mathbb{Z}(\sqrt{21}) = \{m + n\sqrt{21} \mid m, n \in \mathbb{Z}\}$  and  $F = \mathbb{Q}(\sqrt{21})$ , the field of fractions of D. Show the following:
  - (a)  $x^2 x 5$  is irreducible in D[x] but not in F[x].
  - (b) D is not a unique factorization domain.
- 84. Let  $D = \mathbb{Z}(\sqrt{-11}) = \{m + n\sqrt{-11} \mid m, n \in \mathbb{Z}\}$  and  $F = \mathbb{Q}(\sqrt{-11})$  its field of fractions. Show the following:
  - (a)  $x^2 x + 3$  is irreducible in D[x] but not in F[x].
  - (b) D is not a unique factorization domain.
- 85. Let  $D = \mathbb{Z}(\sqrt{13}) = \{m + n\sqrt{13} \mid m, n \in \mathbb{Z}\}$  and  $F = \mathbb{Q}(\sqrt{13})$  its field of fractions. Show the following:
  - (a)  $x^2 + 3x 1$  is irreducible in D[x] but not in F[x].
  - (b) D is not a unique factorization domain.
- 86. Let D be an integral domain and F a subring of D that is a field. Show that if each element of D is algebraic over F, then D is a field.
- 87. Let R be an integral domain containing the subfield F and assume that R is finite dimensional over F when viewed as a vector space over F. Prove that R is a field.
- 88. Let D be an integral domain.
  - (a) For  $a, b \in D$  define a greatest common divisor of a and b.
  - (b) For  $x \in D$  denote  $(x) = \{dx \mid d \in D\}$ . Prove that if (a) + (b) = (d), then d is a greatest common divisor of a and b.
- 89. Let D be a principal ideal domain.
  - (a) For  $a, b \in D$ , define a least common multiple of a and b.
  - (b) Show that  $d \in D$  is a least common multiple of a and b if and only if  $(a) \cap (b) = (d)$ .
- 90. Let D be a principal ideal domain and let  $a, b \in D$ .
  - (a) Show that there is an element  $d \in D$  that satisfies the properties i. d|a and d|b and
    - ii. if e|a and e|b then e|d.
  - (b) Show that there is an element  $m \in D$  that satisfies the properties i. a|m and b|m and
    - ii. if a|e and b|e then m|e.
- 91. Let R be a principal ideal domain. Show that if (a) is a nonzero ideal in R, then there are only finitely many ideals in R containing (a).

- 92. Let D be a unique factorization domain and F its field of fractions. Prove that if d is an irreducible element in D, then there is no  $x \in F$  such that  $x^2 = d$ .
- 93. Let D be a Euclidean domain. Prove that every non-zero prime ideal is a maximal ideal.
- 94. Let  $\pi$  be an irreducible element of a principal ideal domain R. Prove that  $\pi$  is a prime element (that is,  $\pi \mid ab$  implies  $\pi \mid a$  or  $\pi \mid b$ ).
- 95. Let D with  $\varphi : D \{0\} \to \mathbb{N}$  be a Euclidean domain. Suppose  $\varphi(a + b) \leq \max\{\varphi(a), \varphi(b)\}$  for all  $a, b \in D$ . Prove that D is either a field or isomorphic to a polynomial ring over a field.
- 96. Let D be an integral domain and F its field of fractions. Show that if g is an isomorphism of D onto itself, then there is a unique isomorphism h of F onto F such that h(d) = g(d) for all d in D.
- 97. Let D be a unique factorization domain such that if p and q are irreducible elements of D, then p and q are associates. Show that if A and B are ideals of D, then either  $A \subseteq B$  or  $B \subseteq A$ .
- 98. Let D be a unique factorization domain and p a fixed irreducible element of D such that if q is any irreducible element of D, then q is an associate of p. Show the following.
  - (a) If d is a nonzero element of D, then d is uniquely expressible in the form  $up^n$ , where u is a unit of D and n is a non-negative integer.
  - (b) D is a Euclidean domain.
- 99. Prove that  $\mathbb{Z}[\sqrt{-2}] = \{a + b\sqrt{-2} \mid a, b \in \mathbb{Z}\}$  is a Euclidean domain.
- 100. Show that the ring  $\mathbb{Z}[i]$  of Gaussian integers is a Euclidean ring and compute the greatest common divisor of 5 + i and 13 using the Euclidean algorithm.

#### **Polynomial Rings**

- 101. Show that the polynomial  $f(x) = x^4 + 5x^2 + 3x + 2$  is irreducible over the field of rational numbers.
- 102. Let D be an integral domain and D[x] the polynomial ring over D. Suppose  $\varphi : D[x] \to D[x]$  is an isomorphism such that  $\varphi(d) = d$  for all  $d \in D$ . Show that  $\varphi(x) = ax + b$  for some  $a, b \in D$  and that a is a unit of D.
- 103. Let  $f(x) = a_0 + a_1x + \cdots + a_kx^k + \cdots + a_nx^n \in \mathbb{Z}[x]$  and p a prime such that  $p|a_i$  for  $i = 1, \ldots, k 1, p \nmid a_k, p \nmid a_n$ , and  $p^2 \nmid a_0$ . Show that f(x) has an irreducible factor in  $\mathbb{Z}[x]$  of degree at least k.
- 104. Let D be an integral domain and D[x] the polynomial ring over D in the indeterminate x. Show that if every nonzero prime ideal of D[x] is a maximal ideal, then D is a field.
- 105. Let R be a commutative ring with 1 and let  $f(x) \in R[x]$  be nilpotent. Show that the coefficients of f are nilpotent.
- 106. Show that if R is an integral domain and f(x) is a unit in the polynomial ring R[x], then f(x) is in R.

- 107. Let D be a unique factorization domain and F its field of fractions. Prove that if f(x) is a monic polynomial in D[x] and  $\alpha \in F$  is a root of f, then  $\alpha \in D$ .
- (a) Show that x<sup>4</sup> + x<sup>3</sup> + x<sup>2</sup> + x + 1 is irreducible in Z<sub>3</sub>[x].
  (b) Show that x<sup>4</sup> + 1 is not irreducible in Z<sub>3</sub>[x].
- 109. Let F[x, y] be the polynomial ring over a field F in two indeterminates x, y. Show that the ideal generated by  $\{x, y\}$  is not a principal ideal.
- 110. Let F be a field. Prove that the polynomial ring F[x] is a PID and that F[x, y] is not a PID.
- 111. Let D be an integral domain and let c be an irreducible element in D. Show that the ideal (x, c) generated by x and c in the polynomial ring D[x] is not a principal ideal.
- 112. [CORRECTED] Show that if R is a commutative ring with 1 that is not a field, then R[x] is not a principal ideal domain.
- 113. (a) Let  $\mathbb{Z}\begin{bmatrix} \frac{1}{2} \end{bmatrix} = \{\frac{a}{2^n} \mid a, n \in \mathbb{Z}, n \ge 0\}$ , the smallest subring of  $\mathbb{Q}$  containing  $\mathbb{Z}$  and  $\frac{1}{2}$ . Let (2x - 1) be the ideal of  $\mathbb{Z}[x]$  generated by the polynomial 2x - 1. Show that  $\mathbb{Z}[x]/(2x - 1) \cong \mathbb{Z}\begin{bmatrix} \frac{1}{2} \end{bmatrix}$ .
  - (b) Find an ideal I of  $\mathbb{Z}[x]$  such that  $(2x-1) \subsetneq I \subsetneq \mathbb{Z}[x]$ .

# **Non-commutative Rings**

- 114. Let R be a ring with identity such that the identity map is the only ring automorphism of R. Prove that the set N of all nilpotent elements of R is an ideal of R.
- 115. Let p be a prime. A ring S is called a p-ring if the characteristic of S is a power of p. Show that if R is a ring with identity of finite characteristic, then R is isomorphic to a finite direct product of p-rings for distinct primes.
- 116. Let R be a ring.
  - (a) Show that there is a unique smallest (with respect to inclusion) ideal A such that R/A is a commutative ring.
  - (b) Give an example of a ring R such that for every proper ideal I, R/I is not commutative. Verify your example.
  - (c) For the ring  $R = \left\{ \begin{bmatrix} a & b \\ 0 & c \end{bmatrix} \mid a, b, c \in \mathbb{Z} \right\}$  with the usual matrix operations, find the ideal A of part (a).
- 117. If R is any ring with identity, let J(R) denote the Jacobson radical of R. Show that if e is any idempotent of R, then J(eRe) = eJ(R)e.
- 118. If n is a positive integer and F is any field, let  $M_n(F)$  denote the ring of  $n \times n$  matrices with entries in F. Prove that  $M_n(F)$  is a simple ring. Equivalently,  $\operatorname{End}_F(V)$  is a simple ring if V is a finite dimensional vector space over F.

- 119. A ring R is *nilpotent-free* if  $a^n = 0$  for  $a \in R$  and some positive integer n implies a = 0.
  - (a) Suppose there is an ideal I such that R/I is nilpotent-free. Show there is a unique smallest (with respect to inclusion) ideal A such that R/A is nilpotent-free.
  - (b) Give an example of a ring R such that for every proper ideal I, R/I is not nilpotent-free. Verify your example.
  - (c) Show that if R is a commutative ring with identity, then there is a proper ideal I of R such that R/I is nilpotent-free, and find the ideal A of part (a).

#### Local Rings, Localization, Rings of Fractions

- 120. Let R be an integral domain. Construct the field of fractions F of R by defining the set F and the two binary operations, and show that the two operations are well-defined. Show that F has a multiplicative identity element and that every nonzero element of F has a multiplicative inverse.
- 121. A *local* ring is a commutative ring with 1 that has a unique maximal ideal. Show that a ring R is local if and only if the set of non-units in R is an ideal.
- 122. Let R be a commutative ring with  $1 \neq 0$  in which the set of nonunits is closed under addition. Prove that R is local, i.e., has a unique maximal ideal.
- 123. Let D be an integral domain and F its field of fractions. Let P be a prime ideal in D and  $D_P = \{ab^{-1} \mid a, b \in D, b \notin P\} \subseteq F$ . Show that  $D_P$  has a unique maximal ideal.
- 124. Let R be a commutative ring with identity and M a maximal ideal of R. Let  $R_M$  be the ring of quotients of R with respect to the multiplicative set  $R M = \{s \in R \mid s \notin M\}$ . Show the following.
  - (a)  $M_M = \{ \frac{a}{s} \mid a \in M, s \notin M \}$  is the unique maximal ideal of  $R_M$ .
  - (b) The fields R/M and  $R_M/M_M$  are isomorphic.
- 125. Let R be an integral domain, S a multiplicative set, and let  $S^{-1}R = \{\frac{r}{s} \mid r \in R, s \in S\}$  (contained in the field of fractions of R). Show that if P is a prime ideal of R, then  $S^{-1}P$  is either a prime ideal of  $S^{-1}R$  or else equals  $S^{-1}R$ .
- 126. Let R be a commutative ring with identity and P a prime ideal of R. Let  $R_P$  be the ring of quotients of R with respect to the set  $R P = \{s \in R \mid s \notin P\}$ . Show that  $R_P/P_P$  is the field of fractions of the integral domain R/P.
- 127. Let D be an integral domain and F its field of fractions. Denote by  $\mathcal{M}$  the set of all maximal ideals of D. For  $M \in \mathcal{M}$ , let  $D_M = \{\frac{a}{s} \mid a, s \in D, s \notin M\} \subset F$ . Show that  $\bigcap_{M \in \mathcal{M}} D_M = D$ .
- 128. Let R be a commutative ring with 1 and D a multiplicative subset of R containing 1. Let J be an ideal in the ring of fractions  $D^{-1}R$  and let

$$I = \{ a \in R \mid \frac{a}{d} \in J \text{ for some } d \in D \}.$$

Show that I is an ideal of R.

129. Let D be a principal ideal domain and let P be a non-zero prime ideal. Show that  $D_P$ , the localization of D at P, is a principal ideal domain and has a unique irreducible element, up to associates.

## **Chains and Chain Conditions**

- 130. Let R be a commutative ring with identity. Prove that any non-empty set of prime ideals of R contains maximal *and* minimal elements.
- 131. Let R be a commutative ring with 1. We say R satisfies the ascending chain condition if whenever  $I_1 \subseteq I_2 \subseteq I_3 \subseteq \cdots$  is an ascending chain of ideals, there is an integer N such that  $I_k = I_N$  for all  $k \ge N$ . Show that R satisfies the ascending chain condition if and only if every ideal of R is finitely generated.
- 132. **[NEW]** Define Noetherian ring and prove that if R is Noetherian, then R[x] is Noetherian.
- 133. Let R be a commutative Noetherian ring with identity. Prove that there are only finitely many *minimal* prime ideals of R.
- 134. [NEW] Let R be a commutative Noetherian ring in which every 2-generated ideal is principal. Prove that R is a Principal Ideal Domain.
- 135. Let R be a commutative Noetherian ring with identity and let I be an ideal in R. Let J = Rad(I). Prove that there exists a positive integer n such that  $j^n \in I$  for all  $j \in J$ .
- 136. Let R be a commutative Noetherian domain with identity. Prove that every nonzero ideal of R contains a product of nonzero *prime* ideals of R.
- 137. Let R be a ring satisfying the descending chain condition on right ideals. If J(R) denotes the Jacobson radical of R, prove that J(R) is nilpotent.
- 138. Show that if R is a commutative Noetherian ring with identity, then the polynomial ring R[x] is also Noetherian.
- 139. Let P be a nonzero prime ideal of the commutative Noetherian domain R. Assume P is principal. Prove that there does not exist a prime ideal Q satisfying (0) < Q < P.
- 140. Let R be a commutative Noetherian ring. Prove that every nonzero ideal A of R contains a product of prime ideals (not necessarily distinct) each of which contains A.
- 141. Let R be a commutative ring with 1 and let M be an R-module that is not Artinian (Noetherian, of finite composition length). Let  $\mathcal{I}$  be the set of ideals I of R such that there exists an R-submodule N of M with the property that N/NI is not Artinian (Noetherian, of finite composition length, respectively). Show that if  $A \in \mathcal{I}$  is a maximal element of  $\mathcal{I}$ , then A is a prime ideal of R.